

Longitudinal Momentum Fraction X_L for Two High P_t Protons in $pp \rightarrow ppX$ Reaction

D. Zhalov^e, S. Heppelmann^e, J. Alster^a, G. Asryan^{c,b},
Y. Averichev^h, D. Barton^c, V. Baturin^{e,d}, N. Bukhtoyarova^{c,d},
A. Carroll^c, T. Kawabata^f, A. Leksanov^e, Y. Makdisi^c, A. Malki^a,
E. Minina^e, I. Navon^a, H. Nicholson^g, A. Ogawa^e, Yu. Panebratsev^h,
E. Piassetzky^a, A. Schetkovsky^{e,d}, S. Shimanskiy^h, A. Tangⁱ,
J.W. Watsonⁱ, H. Yoshida^f

^a*School of Physics and Astronomy, Sackler Faculty of Exact Sciences, Tel Aviv University,
Ramat Aviv 69978, Israel*

^b*Yerevan Physics Institute, Yerevan 375036, Armenia*

^c*Collider-Accelerator Department, Brookhaven National Laboratory, Upton, NY 11973, USA*

^d*Petersburg Nuclear Physics Institute, Gatchina, St. Petersburg 188350, Russia*

^e*Physics Department, Pennsylvania State University, University Park, PA 16801, USA*

^f*Department of Physics, Kyoto University, Sakyo-ku, Kyoto, 606-8502, Japan*

^g*Department of Physics, Mount Holyoke College, South Hadley, MA 01075, USA*

^h*J.I.N.R., Dubna, 141980, Russia*

ⁱ*Department of Physics, Kent State University, Kent, OH 44242, USA*

Abstract. We present an analysis of new data from Experiment E850 at BNL. We have characterized the inclusive cross section near the endpoint for pp exclusive scattering in Hydrogen and in Carbon with incident beam energy of 6 GeV. We select events with a pair of back-to-back hadrons at large transverse momentum. These cross sections are parameterized with a form $\frac{d\sigma}{dX_L} \sim (1 - X_L)^p$, where X_L is the ratio of the longitudinal momentum of the observed pair to the total incident beam momentum. Small value of p may suggest that the number of partons participating in the reaction is large and reaction has a strong dependence on the center-of-mass energy. We also discuss nuclear effects observed in our kinematic region.

There are two main questions which can be addressed in the study of the Longitudinal Momentum Fraction carried by a pair of large P_t final state particles in the $pp \rightarrow ppX$ reactions. We learn first something about the number of partons involved in the interaction. Concluding, as we will, that the number of partons involved is large, we discuss a mechanism for picking up a momentum boost from the spectator nucleus in events where the observed momentum fraction is equal and larger than

the incoming momentum.

We defined the Longitudinal Momentum Fraction variable X_L as:

$$X_L = \frac{P_{L_1} + P_{L_2}}{P_{L_{inc}}}. \quad (1)$$

Here $P_{L_{inc}}$ is the momentum of the beam coming along the Z-axis, P_{L_1} and P_{L_2} are the longitudinal components of the two detected out-going particles having large transverse momenta. We have argued that the two detected out-going particles in the $pp \rightarrow ppX$ reaction are protons [1]. We then expect the shape of the cross section, as X_L approaches unity, to reflect the number of partons participating in the collision. If the cross section is parameterized as

$$\frac{d\sigma}{dX_L} \sim (1 - X_L)^p \quad (2)$$

then a large p is expected for single parton collisions, reflecting the endpoint behavior of structure functions and fragmentation functions [2]. As the number of participating partons increases, we expect the power p to be reduced.

In the experiment discussed here, we measure p for selected classes of two track events. Our result, that p is in the range of 1-2, indicates that many partons are participating in the hard process. It can be expected that the cross section may be falling very rapidly with center-of-mass (cm) energy for these events, perhaps similar to the pp elastic energy dependence.

The cm energy dependence of the underlying $pp \rightarrow ppX$ cross section is affected by the number of partons participating in the interaction. The underlying cross section falls with energy as a power of energy, and that power increases as the number of partons increases. It is known that for exclusive scattering, the cross section has a dependence on the cm energy given by (s is a Mandelstam variable) $\sigma \sim s^{-m}$, $m = (n_1 + n_2 + n_3 + n_4 - 2)$ for a process with $n_1 + n_2$ partons $\rightarrow n_3 + n_4$ partons [3].

With this observation about the nature of the selected $pp \rightarrow ppX$ events, we can further conclude that interesting nuclear effects would also be expected. The nucleus can change the cm energy of the incident proton and the nuclear proton by virtue of nuclear fermi momentum. If the nucleus contributes a target proton which is moving in the direction of the beam particle, the cm energy is reduced. Because the increase in cross section associated with a reduction in cm energy is very large, we can expect a strong bias for the nucleus to contribute momentum to the scattering process in the forward direction. The typical magnitude of this nuclear contribution to the final state momentum may be much larger than typical fermi momentum seen in other processes.

Since the nucleus tends to add positive momentum to the two final state particles, in some sense, the stopping power for these events is apparently negative. When one compares the same $pp \rightarrow ppX$ processes measured in nuclear matter with that measured in Hydrogen, the effect of the nucleus is to move strength in the X_L

distribution to larger X_L . If a ratio, like a transparency ratio of the cross section in nuclear matter to that measured in Hydrogen is calculated, the ratio will be artificially enhanced.

The data used for this analysis was collected on two targets ^{12}C and CH_2 using the EVA Spectrometer (E850 Exp., BNL, AGS). The CH_2 target enabled us to extract cross sections for free proton (Hydrogen) scattering. The acceptance of the detector for two high P_t tracks in the r-z plane was centered around the opening angle for elastic $\text{pp} \rightarrow \text{pp}$ scattering at 90° in the cm frame [4]. Charged particle acceptance in the azimuthal angle was nearly 2π .

In Fig. 1, we present the X_L distribution for Carbon and Hydrogen. In one case we applied a 'semi-elastic' cut (two and only two charged tracks in our charged track acceptance) which allowed us to look at the X_L distribution for $\text{pp} \rightarrow \text{pp}$ events with a background of $\text{pp} \rightarrow \text{pp} + X^0$ and $\text{pp} \rightarrow \text{pp} + X^{\text{charged}}$. The quasi-elastic signals have been extensively analyzed elsewhere [4,5].

Secondly, we select a more inclusive data set made up of events with two high P_t tracks but any number of additional lower momentum charged tracks in the $\text{pp} \rightarrow \text{ppX}$ process.

We have fitted X_L distribution with a form (here A and p are the fit constants):

$$\frac{d\sigma}{dX_L} = A * (1 - X_L)^p + \text{Gaussian} \quad (3)$$

The polynomial part of the fit was restricted for $X_L < 1$ while the Gaussian is included to account for the quasi-elastic contributions near $X_L \simeq 1$.

The peak corresponding to the Gaussian part of the fit is actually shifted for Carbon events from $X_L = 1$ towards larger X_L because of the s^{-10} dependence of the $\text{pp} \rightarrow \text{pp}$ reaction's cross section.

We have determined the power of the polynomial in the fit to be in the range 1.1-1.8 depending on the tightness of the cuts around the region of the elastic kinematics. According to our interpretation, we conclude that the number of quarks participating in the process is large. Consequently, the power m of s^{-m} dependence of the cross section would be expected to be large as well.

In Fig. 1, the effective ratio for scattering in Carbon vs Hydrogen is shown. While the ratio is well in excess of Glauber levels, our interpretation is that the Carbon distribution is enhanced due to the strong energy dependence of the cross section.

A recent study of neutron correlations from this experiment (E850 collaboration at BNL) has shown an anomalously large yield of neutrons going into the backward hemisphere in $\text{pp} \rightarrow \text{ppX}$ reactions compared to the yields reported by other experiments [1,6]. This result was observed with a data sample similar to the one described above. A possible interpretation of this phenomenon is that strong s-dependence of the cross-section for this class of events may select events with large positive fermi momentum. The nucleus must balance this momentum by some kind of recoil in the backward direction. The idea that most of the recoil momentum is

absorbed by a single recoiling neutron leads to the prediction of enhanced backward neutron production [7]. Since a large yield of backward neutrons has been observed in the $pp \rightarrow ppX$ data set described here, the picture seems to have merit.

Keeping this in mind, we looked at the X_L distribution for Carbon and Hydrogen (Fig. 1). In the case of exclusive events with two high P_t particles, we can clearly see the shift in the position of the Gaussian peak in the Carbon sample. This effect is known to be caused by the s^{-10} dependence of the cross section. We have compared the more inclusive data set distributions for Carbon and Hydrogen for $pp \rightarrow ppX$ and interpreted the enhanced ratio as evidence for a similar shift in the Carbon distribution with respect to the Hydrogen for this more inclusive sample as well.




FIGURE 1. LEFT: X_L distributions for Carbon nucleus and 6*(Hydrogen) with two and only two tracks cut. RIGHT: Ratio of the Carbon signal to the 6*(Hydrogen) signal with two and only two tracks cut.

REFERENCES

1. Malki A. et al, *to be published*.
2. Brodsky S. et al, *Nucl. Phys.* **B441**, 197 (1995).
3. Brodsky S., Farrar G., *Phys. Rev. Lett.* **31**, 1153 (1973).
4. Leksanov A. et al, in *Proceedings of CIPANP2000, Quebec, May 22-28, 2000*.
5. Carroll A. et al, *Phys. Rev. Lett.* **61**, 1698 (1988).
6. Malki A. et al, in *Proceedings of CIPANP2000, Quebec, May 22-28, 2000*.
7. Frankfurt L.L., Strikman M.I., *Phys. Lett.* **B69**, 87 (1977).